

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

QUANTUM

Carfentrazone 400 EC

HERBICIDE

ACTIVE CONSTITUENT: 400 g/L CARFENTRAZONE-ETHYL

SOLVENT: 552 g/L HYDROCARBON LIQUID

GROUP **14** HERBICIDE

For improvement in the control of marshmallow and certain other broadleaf weeds prior to establishment of crops, fallows or forest plantations, in commercial, industrial and public service areas, around agricultural buildings and yards, in tree fruits, nuts, grape vines in tank mixture with knockdown herbicides; control of marshmallow and annual nettles in grass pastures and rough grass/turf areas; control of volunteer cotton seedlings including Roundup Ready cotton and desiccation of cotton re-growth; control of unwanted suckers (cane burning) in *Rubus* spp. and *Ribes* spp. as per the Directions for Use Table.

APVMA Approval No.: 91876/150250

EC Formulation Type
Emulsifiable
Concentrate

**BE PROUDLY
AUSTRALIAN**

Quantum Agrosiences Holdings Pty Ltd | ABN 79 680 792 625
Suite 2, Level 7, 330 Collins Street, Melbourne, Victoria 3000
T: 1300 658 988 | www.quantumag.au

 **QUANTUM
AGROSCIENCES**

DIRECTIONS FOR USE

RESTRAINT: DO NOT apply by aerial application except for cotton desiccation.

SITUATION	WEEDS CONTROLLED	STATE	TIME OF APPLICATION
<p>Prior to sowing winter and summer broadacre and horticultural crops; starting at fallow; and prior to establishment of forestry plantations.</p> <p>To assist in weed control in Commercial, Industrial and Public Service areas, around Agricultural buildings and yards.</p> <p>In young or established Grapevines, Treefruits (including Pome fruit, Stone fruit and Citrus fruit), Tree Nuts and Assorted Tropical and Sub tropical fruits.</p>	<p>Australian Crassula/Stoncrop <i>Crassula</i> spp., Capeweed <i>Arctotheca calendula</i>, Chickweed <i>Stellaria media</i>, Common storksbill (max. 4 leaves) <i>Erodium cicutarium</i>, Doublegee/Spiny emex/Three cornered jack <i>Emex australis</i>, Marshmallow <i>Malva parviflora</i>, Paterson's curse <i>Echium plantagineum</i>, Sub. clover <i>Trifolium subterraneum</i>, Wild radish <i>Raphanus raphanistrum</i></p> <p>Refer also to the product label for the knockdown herbicide used.</p> <p>If one of the above weeds is the dominant weed, and there is no specific rate for it in the knockdown herbicide's label, consult the label's generic annual- weed rate-range.</p> <p>Select from within this range to suit the weed-stage, weed-density, conditions (etc) of your situation.</p>	All States	<p>Apply as a tank mix with Wipe-Out 450, Wipe-Out Accelerate or other glyphosate products;</p> <p>Spray & Sow, Spraynut or other paraquat products.</p>

RATE	CRITICAL COMMENTS
<p>15 - 45 mL/ha plus recommended label rates of knockdown herbicides</p> <p>Spot spray 6 mL/100 L plus recommended label spot spray rates of knockdown herbicides</p>	<p>Addition of Quantum Carfentrazone 400 EC Herbicide to knockdown herbicides will increase the speed at which treated broadleaved weeds in general develop visible symptoms (compared to results achieved with knockdown herbicides applied alone) and may improve final control of broadleaved weeds including certain hard-to-kill weeds, marshmallow in particular. The use of higher rates and full soil disturbance in cropping situations may improve control of marshmallow in particular.</p> <p>Use the lower rates on younger plants or plants growing under good conditions and the higher rates on older plants or plants growing under less optimum conditions.</p> <p>The lower rate may only provide suppression of capeweed, wild radish, common storksbill and doublegee under poor growing conditions. Common storksbill should be no larger than 4 leaf at spraying, recently germinated and not under stress – older or stressed plants may not be adequately controlled.</p> <p>Application to hardened weeds or drought stressed weeds especially under summer conditions may cause only localised injury to weed foliage which may not enhance final weed control. Mallow growing and sprayed in the summer is especially prone to drought stress and may either not show symptoms typical of Quantum Carfentrazone 400 EC Herbicide or may re-grow following treatment although plants did not appear very stressed at application.</p> <p>Apply only as a tank mix with recommended rates of knockdown herbicides. Refer to the appropriate label for weed sizes and follow all label directions. Addition of a paraffinic oil adjuvant at 0.5% may be beneficial when applying Quantum Carfentrazone 400 EC Herbicide with a glyphosate herbicide.</p> <p>To ensure uptake of Quantum Carfentrazone 400 EC Herbicide DO NOT sow crops for at least 1 hour after application. Always refer to the appropriate companion product label in case a longer re-crop sowing period is required.</p> <p>When using Quantum Carfentrazone 400 EC Herbicide as a spot spray, apply in sufficient water (minimum 500 L/ha) to thoroughly wet all weed foliage to the point of run-off. Addition of standard rates of a non-ionic surfactant may improve weed control.</p>

SITUATION	WEEDS CONTROLLED	STATE	TIME OF APPLICATION
Prior to sowing summer broadacre and horticultural crops; starting a fallow	Volunteer cotton seedlings, including Roundup Ready* varieties	NSW, Qld, WA only	Apply to seedlings at 2 - 6 leaf stage
Grass pastures; Rough grass/turf areas	Marshmallow <i>Malva parviflora</i> , Annual (stinging) nettles <i>Urtica urens</i>	All States	Apply to seedlings at 2 - 10 leaf stage
Cotton	Desiccation of re-growth	Qld, NSW & WA only	Apply to re-growth following defoliation

RATE	CRITICAL COMMENTS
<p>Roundup Ready cotton only: 45 - 60 mL/ha plus a paraffinic oil adjuvant or Hasten* 1%</p> <p>or</p> <p>45 - 60 mL/ha plus Wipe-Out 450, Wipe-Out Accelerate plus a paraffinic oil adjuvant or Hasten 0.5%</p> <p>Conventional cotton only: 30 - 45 mL/ha plus recommended rates of knockdown herbicides</p>	<p>Apply Quantum Carfentrazone 400 EC Herbicide in a minimum spray volume of 80 L/ha to ensure effective coverage. Use of lower spray volumes may reduce weed control. Cotton should be growing well at application; application to plants growing under heat and moisture stress may reduce the level of control.</p> <p>To broaden the weed spectrum Quantum Carfentrazone 400 EC Herbicide may be tank mixed with the recommended rate of a knockdown herbicide (Wipe-Out 450, Wipe-Out Accelerate or other glyphosate products, Spraytop or other paraquat products, Spray & Sow). When using a tank mix with glyphosate for control of Roundup Ready seedlings the higher rate range of Quantum Carfentrazone 400 EC Herbicide is required as well as a paraffinic oil adjuvant or Hasten at 0.5%.</p> <p>When using a tank mix with glyphosate for control of Conventional Cotton seedlings the addition of standard rates of a non-ionic surfactant or 0.5% of an oil adjuvant such as a paraffinic oil or Hasten will maximise control.</p>
<p>15 - 45 mL/ha plus a paraffinic oil adjuvant or Hasten 0.5%</p> <p>15 - 45 mL/ha plus recommended rates of 2,4-D</p>	<p>Use the lower rates on younger plants or plants growing under good conditions and the higher rates on older plants or plants growing under less optimum conditions. These rates may only provide suppression of marshmallow.</p> <p>To improve the control of marshmallow Quantum Carfentrazone 400 EC Herbicide should be tank mixed with one of the following: 500 mL/ha of an 800 g/L ethyl ester of 2,4-D; 650 mL/ha of a 625 g/L amine of 2,4-D - activity of this mix will be improved with the addition of a paraffinic oil adjuvant or Hasten at 0.5%.</p>
<p>50 - 60 mL/ha plus crop oil concentrates or blended oil/surfactant</p>	<p>Use Quantum Carfentrazone 400 EC Herbicide to desiccate re-growth which occurs following the defoliation program.</p> <p>Apply with crop oil concentrates or blended oil/surfactant adjuvants as this may result in a greater reduction of green shoot.</p> <p>Use the higher rate of Quantum Carfentrazone 400 EC Herbicide when re-growth is vigorous or when there are more than 20 re-growth leaves of any size, per plant.</p> <p>Ensure that spray equipment is adjusted to provide thorough coverage of foliage to maximise product performance. Use a minimum of 100 L/ha for ground application and 30 L/ha for aerial application.</p>

SITUATION	WEEDS CONTROLLED	STATE	TIME OF APPLICATION
<i>Rubus</i> spp. (including raspberries and blackberries) and <i>Ribes</i> spp. (including currants and gooseberries)	Control of unwanted suckers (cane burning)	All States and Territories	Apply to suckers less than 300 mm long, before any lignification occurs.

RATE	CRITICAL COMMENTS
45 mL/100 L	<p>Apply to suckers less than 300 mm long, before lignification occurs.</p> <p>Ensure complete coverage of suckers and spray to point of run-off.</p> <p>A minimum spray volume of 50 L per kilometre of row (both sides sprayed) is recommended.</p> <p>Avoid spray drift onto desirable green stems, foliage, fruit or flowers.</p> <p>Use hooded or shielded sprayers when applying treatment in close proximity to seedlings or newly transplanted bushes, or when planting distance between bushes and distances between rows are minimal.</p> <p>WARNING: This product is highly toxic to foliage and green stems of broadleaf plants, and incidental spray drift onto desirable green stems, foliage, fruit or flowers may result in necrosis.</p> <p>The spraying of target emergent suckers is not known to affect the health of <i>Rubus</i> or <i>Ribes</i> bush varieties; however, it is recommended to treat a small area of suckers in close proximity to <i>Rubus</i> or <i>Ribes</i> bushes to ascertain any response before treating the whole crop.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED FOR QUANTUM CARFENTRAZONE 400 EC HERBICIDE WHEN USED AS DIRECTED.

HOWEVER, REFER ALSO TO THE WITHHOLDING PERIOD OF PRODUCT/S MIXED WITH QUANTUM CARFENTRAZONE 400 EC HERBICIDE.

GRAZING: DO NOT GRAZE TREATED AREAS OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER APPLICATION.

COTTON PROCESSING: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

GRAZING/COTTON TRASH: DO NOT GRAZE TREATED AREAS OR FEED COTTON TRASH TO LIVESTOCK.

EXPORT SLAUGHTER INTERVAL (ESI)

An export slaughter interval is not required when used as directed.

GENERAL INSTRUCTIONS

Quantum Carfentrazone 400 EC Herbicide is an emulsifiable concentrate formulation and is to be mixed with water and applied as per the labelled Directions for Use.

Quantum Carfentrazone 400 EC Herbicide is a post-emergence herbicide to be added to knockdown herbicides to improve the control of certain broadleaf weeds including marshmallow prior to the establishment of fallows, winter and summer broadacre and horticultural crops, marshmallow control in grass pastures, in commercial, industrial and public service areas, around agricultural buildings, yards and other farm situations or in treefruits, nuts and grapevines. Quantum Carfentrazone 400 EC Herbicide can be used alone with a suitable adjuvant for control of volunteer cotton seedlings including Roundup Ready cotton. It can also be used for desiccation of regrowth in cotton.

Quantum Carfentrazone 400 EC Herbicide is a fast-acting contact herbicide and aids in control of weeds through a process of membrane disruption. The foliar uptake of Quantum Carfentrazone 400 EC Herbicide is rapid and plant desiccation can occur within 4 days of application. Application of Quantum Carfentrazone 400 EC Herbicide should target small actively growing weeds. Subsequent germinations will not be controlled.

SYMPTOMS

Quantum Carfentrazone 400 EC Herbicide is rapidly absorbed through the foliage of plants. Within a few hours following application, the foliage of susceptible weeds shows signs of desiccation, and in subsequent days necrosis and death of the plant.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions and cultural practices may affect the activity of Quantum Carfentrazone 400 EC Herbicide. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicidal symptoms is delayed, and weeds hardened off by drought are less susceptible to Quantum Carfentrazone 400 EC Herbicide.

COMPATIBILITY

For most uses as per the Directions for Use Quantum Carfentrazone 400 EC Herbicide should always be tank mixed with formulations of the knockdown herbicides including Wipe-Out 450, Wipe-Out Accelerate, Credit*/Bonus* or other products based on glyphosate, Spray & Sow, Spraytop or other knockdown herbicides based on paraquat and Exonerate. Quantum Carfentrazone 400 EC Herbicide is also compatible with partner herbicides commonly used with knockdown herbicides including Farmozine, atrazine SC, Banvel* 200, 2,4-D amine, 2,4-D ester, Lonestar® (triasulfuron), pendimethalin, Simanex® 900 WG, simazine SC and Trilogy® (trifluralin). This compatibility claim is

restricted to a three-way mix of Quantum Carfentrazone 400 EC Herbicide with any one of the above partner herbicides plus a knockdown herbicide (provided the knockdown herbicide label includes a claim of compatibility with that partner herbicide). Quantum Carfentrazone 400 EC Herbicide is compatible with non-ionic surfactants (wetting agents) such as Wetspray® 1000, Shirwet* 600 and with oil adjuvants including paraffinic oil and Hasten.

TIMING

Pre-crop, Volunteer cotton and Marshmallow in grass pastures

Application should be made to small, actively growing weeds less than 6 to 8 leaf in stage. As Quantum Carfentrazone 400 EC Herbicide is a contact herbicide, best control is achieved when weeds are exposed and not shielded by other weeds and or stubble.

Cotton

If cotton re-growth occurs after the initial defoliation program, apply Quantum Carfentrazone 400 EC Herbicide to desiccate re-growth prior to harvest.

MIXING

Add half the required volume of water to spray tank and start agitation. Add any partner SC or WG herbicide next if it should be added before an EC or knockdown herbicide. Add the measured amount of Quantum Carfentrazone 400 EC Herbicide next, followed by the knockdown herbicide (if required). Add balance of water to tank and add any adjuvant or additive if recommended for use with the knockdown herbicide or for use in grapevines or as suggested in the Critical Comments. Maintain good agitation at all times until spraying is completed. DO NOT store spray mixture overnight.

APPLICATION

DO NOT apply by aerial application except for cotton desiccation.

Ground sprayers

Pre-crop, Volunteer cotton and Marshmallow in grass pastures

Apply Quantum Carfentrazone 400 EC Herbicide as a broadcast application using a conventional boom sprayer with either mechanical or by-pass agitation. Use single orifice flat fan nozzles such as Spraying Systems TeeJet* 11001, 110015, 11002 or equivalent sizes from other manufacturers or Spraying Systems TwinJet* twin flat spray tips TJ60-11002, TJ60-11003 or TJ60-11004 or equivalent sizes from other manufacturers.

DO NOT use 110-03 or bigger single orifice nozzles or TJ11006 or bigger twin orifice nozzles with Quantum Carfentrazone 400 EC Herbicide. DO NOT use floodjet, low drift or air induction nozzles, boomless jets or misters or controlled droplet application equipment.

Spray equipment should be properly calibrated to ensure correct and uniform application. Use a spray volume of 50 to 150 litres per hectare (minimum 80 L/ha for volunteer cotton). Experience has shown that increasing spray volumes can improve weed control. Use the lowest pressure and boom height which provides uniform coverage. Use the higher volume if weed infestation is heavy and/or tall.

Cotton

When desiccation cotton regrowth, use sufficient water to obtain complete coverage of all leaves, i.e. at least 100 litres of water per hectare. Good coverage should be obtained with three or more nozzles per row depending upon crop height and canopy density.

Knapsack, Pneumatic, Handgun sprayers

Apply Quantum Carfentrazone 400 EC Herbicide at the recommended rate of 1 mL per 10 litres (10 mL per 100 litres) in conjunction with labelled handgun rates of a knockdown herbicide in sufficient water to adequately and uniformly wet the foliage of the weeds being sprayed. A spray volume of 500 L to 1000 L will usually be sufficient but higher volumes may be required for dense taller vegetation. The best application conditions are when soil is moist, weather fine and rain unlikely within one hour or as specified for the knockdown herbicide.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions and/or cultural practices may affect the activity of Quantum Carfentrazone 400 EC Herbicide. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicidal symptoms is delayed, and weeds hardened off by drought are less susceptible to Quantum Carfentrazone 400 EC Herbicide.

Aerial Application (for Cotton Desiccation Only)

Apply by fixed wing aircraft in a minimum of 30 L/ha water and ensure thorough coverage. DO NOT exaggerate swath width.

SPRAYER CLEAN OUT - AFTER THE USE OF Quantum Carfentrazone 400 EC Herbicide

Thoroughly clean all spray equipment using the following procedure when you have finished spraying highly active materials such as carfentrazone-ethyl. In addition to the following procedure, ensure proper equipment clean-out for any other products mixed with Quantum Carfentrazone 400 EC Herbicide as specified on the other product labels.

IMPORTANT

More complete cleaning can be achieved if the spray equipment is cleaned immediately following each use.

Mix only as much herbicide spray solution as needed at a time.

DO NOT store the sprayer for any extended period of time, especially overnight, with

Quantum Carfentrazone 400 EC Herbicide spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

Preparation of the Cleaning Solution: Prepare a spray equipment cleaning solution by mixing an alkaline detergent e.g. "OMO**" or "SPREE**" at a rate of 100 g for every 100 L of clean water used. Upon completion of applying Quantum Carfentrazone 400 EC Herbicide and before spraying sensitive crops including canola, pulses such as faba beans, lentils, other legumes and cotton:

1. Fill the spray tank with sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles then fill the spray tank to capacity to ensure contact of the solution with all internal surfaces. Let the cleaning solution soak in tank, pump and spray lines overnight.
2. Before further use of the sprayer, operate the spray system for 15 minutes, then completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles.
3. If possible, spray a pesticide requiring an oil adjuvant e.g. Pentagon® & a paraffinic oil adjuvant onto cereals as a further means of removing possible residues of Quantum Carfentrazone 400 EC Herbicide before spraying sensitive crops.
4. Immediately prior to commencement of spraying a sensitive crop, purge the boom lines by operating the spray system onto a fence line or waste area for sufficient time to remove any solution that has been residing in the spray lines. This is also recommended for subsequent tank loads or if the sprayer has been left standing for a period of time containing spray solution.
5. If storing equipment for more than 48 hours, preferred practice is to clean spray equipment as outlined above allowing to soak overnight, drain and flush with fresh water and leave fresh water in the spray tank, hoses, and spray booms until next use. This water must be drained from the spray boom and lines and flushed out with clean water before beginning any application to a sensitive crop.

Properly dispose of all cleaning solution and rinsate safely in accordance with Federal, State, and local regulations and guidelines.

DO NOT apply sprayer cleaning solutions or rinsate to sensitive crops. Should small quantities of Quantum Carfentrazone 400 EC Herbicide remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to sensitive crops and other vegetation.

The above method is only effective if the cleaning solution comes into contact with every surface or contact point that may contain even minute carfentrazone-ethyl residues.

CROP PLANT BACK & ROTATION RECOMMENDATIONS

Quantum Carfentrazone 400 EC Herbicide does not provide residual activity, therefore no crop plant back or rotational restrictions apply. However, check the label of any product mixed with Quantum Carfentrazone 400 EC Herbicide to determine any plant back periods or restrictions on use.

RESISTANT WEEDS WARNING

Quantum Carfentrazone 400 EC Herbicide is a member of the Aryl triazolone group of herbicides. Its mode of action is through a process of membrane disruption, which is initiated by the inhibition of the enzyme protoporphyrinogen oxidase. This inhibition interferes with the chlorophyll biosynthetic pathway. For weed resistance management Quantum Carfentrazone 400 EC Herbicide is a Group 14 herbicide. Some naturally occurring weed biotypes resistant to Quantum Carfentrazone 400 EC Herbicide and other Group 14 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Quantum Carfentrazone 400 EC Herbicide or other Group 14 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Quantum Agrosiences Holdings Pty Ltd accepts no liability for any losses that may result from the failure of Quantum Carfentrazone 400 EC Herbicide to control resistant weeds.

PRECAUTIONS

Re-Entry Period: DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spray equipment, which may cause spray drift onto nearby susceptible plants, adjacent crops, or pastures, or onto wetlands, waterbodies or watercourses.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to algae and aquatic plants. DO NOT contaminate streams, rivers or waterways with Quantum Carfentrazone 400 EC Herbicide or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. May irritate the nose and throat. Will irritate the skin and eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. DO NOT inhale vapour or spray mist. When preparing product for use and using the product wear elbow-length chemical resistant gloves. When mixing, loading, and opening the container and preparing spray and using the prepared spray wear cotton overalls over normal clothing buttoned to the neck and wrist and face shield or goggles. If applying by hand wear cotton overalls over normal clothing buttoned to the neck and wrist and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. If in eyes, wash out immediately with water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) which is available from the supplier.

GROUP	14	HERBICIDE
--------------	-----------	------------------

CONDITIONS OF SALE: Quantum Agrosociences Holdings Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Quantum Agrosociences Holdings Pty Ltd's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Quantum Agrosociences Holdings Pty Ltd has any authority to add to or alter these conditions.

Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture: **Combustible liquid. May be fatal if swallowed and enters airways. Very toxic to aquatic life with long-lasting effects.**

Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Avoid release to the environment.



Quantum Agrosociences Holdings Pty Ltd

ABN 79 680 792 625

Suite 2, Level 7, 330 Collins Street,

Melbourne, Victoria 3000

T: 1300 658 988 | www.quantumag.au

APVMA Approval No.: 91876/150250